



TWO-POINT SECOND BOUNDARY VALUE PROBLEM FOR A QUADRATIC SIMPLE SECOND-ORDER DIFFERENTIAL EQUATION SOLVED BY THE BERNOULLI EQUATION

S. T. Qosimova

Teacher, Department of Higher Mathematics, Fergana Polytechnic Institute, Fergana, Uzbekistan

ABSTRACT

This article examines the uniqueness of the solution of the boundary problem for the second regular ordinary differential equation, which is solved in the Bernoulli equation. The uniqueness of the issue is proved by the principle of extremes.

Keywords: Bernoulli equation, unity of a solution, ordinary differential equation, availability of a solution.

Problem Statement

y'' + P1(x)y' + P2(x)y'^2 + P3(x)y'^3 + P4(x)y'^4 = P(x), x in [x0; x1] (1)

equation and

y'(x0) = y0, y(x1) = y1 (2)

find the function y(x) that satisfies the boundary conditions. Here P(x), P1(x), P2(x), P3(x) - given continuous functions.

Theorem. p1(x) - (1) is a special solution of equation (2) that does not satisfy the boundary condition, P3(x) + 4P4(x)p1(x) = 0 and

P1(x) + 2P2(x)p1(x) + 3P3(x)p1^2(x) + 4P4(x)p1^3(x) = Q1(x), y0 - p1(x0) != 0,

Q1(x) != 0, p1'(x) + P1(x)p1(x) + P2(x)p1^2(x) + P3(x)p1^3(x) = P(x)

if the conditions are satisfied, then the problem S3 has a unique solution.

Proof. Using the notation y' = p(x) in Equation (1), we construct the following

equation p'(x) + P1(x)p(x) + P2(x)p^2(x) + P3(x)p^3(x) = P(x) (3)

and condition p(x0) = y0 (4) from the boundary conditions (2).



The resulting {(3), (4)} is in the new case

$$p(x) = p_1(x) + z(x), \tag{5}$$

by performing the substitution and after some elementary simplification we form the

$$z'(x) + (P_1(x) + 2P_2(x)p_1(x) + 3P_3(x)p_1^2(x) + 4P_4(x)p_1^3(x)) \cdot z + (P_2(x) + 3P_3(x)p_1(x) + 6P_4(x)p_1^2(x)) \cdot z^2 + (P_3(x) + 4P_4(x)p_1(x))z^3(x) + P_4(x)z^4(x) = P(x) - [p_1'(x) + P_1(x)p_1(x) + P_2(x)p_1^2(x) + P_3(x)p_1^3(x) + P_4(x)p_1^4(x)]$$

equation. If we apply the conditions of the theorem to this equation

$$z'(x) + Q_1(x)z(x) = -P_4(x)z^4(x) \tag{6}$$

Based on the substitution of the Bernoulli equation in the form Sx and (4) from the boundary condition (5)

$$z(x_0) = y_0 - p_1(x_0) \tag{7}$$

we create the condition. (6) in equation

$$t(x) = \frac{1}{z^3(x)} \tag{8}$$

by carrying out replacement,

$$t'(x) - 3Q_1(x)t(x) = 2P_3(x) \tag{9}$$

while the equation and (7) condition

$$t(x_0) = \frac{1}{[y_0 - p_1(x_0)]^2} \tag{10}$$

we form the condition. As a result, we come to the new {(9), (10)} issue.

Let's assume, {(9),(10)} let the issue have t_1 as well as t_2 solutions [1-9]. In that case

$$t(x) = t_1(x) - t_2(x) \tag{11}$$

function

$$t'(x) - 3Q_1(x)t(x) = 0, \quad x \in [x_0; x_1], \tag{9'}$$

$$t(0) = 0 \tag{10'}$$

will be the solution to a problem.

Suppose that the problem {(9'),(10')} has a solution $t(x) \neq 0, x \in [x_0; x_1]$.



Since the function $t(x)$ is defined and continuous in the segment $[x_0; x_1]$, it reaches a positive maximum (negative minimum) value at some point $x' \in (x_0; x_1]$ of this segment according to Weierstrass's theorem 2.

We assume that the function $t(x)$ should reach a positive maximum (minus minimum) value $(x_0, x_1]$ in half the range. Assuming that $t(x'_0) > 0 (< 0)$ is a positive maximum (minus minimum) value, then $t'(x') = 0$ equality, as well as $t'(x) - 3Q_1(x)t(x) > 0$ inequality is executed [7-12].

This is contrary to (9'). Hence, $t(x)$ the function is (basically) at $\forall x' \in (x_0, x_1]$ ((9') basically)

$$t(x) \equiv 0, \quad \forall x' \in (x_0, x_1] \tag{12}$$

Based on this (11) $t_1(x) = t_2(x)$. And it turns out that the solution of the issue {(9),(10)} is no more than one. All in all, {(9),(10)} the issue is that if he has a solution in the cut $[x_0; x_1]$, It is the only one. {(9), (10)} since the solution of the issue is unique {(1), (2)} the solution of the issue is also unique. Because, {(9),(10)} the issue {(1), (2)} is an equivalent issue.

Availability of Problem Solutions. (9) Using the Bernoulli method to find the general solution of equation

$$t(x) = u(x) \cdot v(x) \tag{13}$$

apparently looking for. Substituting (13) into (9)

$$u'(x) \cdot v(x) + u(x) \cdot [v'(x) - 2Q_1(x) \cdot v(x)] = 2P_3(x)$$

creating equality, hence

$$v(x) = v(x_0) e^{-3 \int_{x_0}^x Q_1(s) ds},$$

$$u(x) = u(x_0) - \frac{3}{v(x_0)} \cdot \int_{x_0}^x P_3(s) e^{3 \int_{x_0}^s Q_1(\tau) d\tau} ds$$

find the functions and solve the general solution of equation (9) based on (13)

$$t(x) = \left[u(x_0)v(x_0) - 3 \cdot \int_{x_0}^x P_3(s) e^{-2 \int_{x_0}^s Q_1(\tau) d\tau} ds \right] \cdot e^{-2 \int_{x_0}^x Q_1(s) ds},$$

$$t(x) = \left[t(x_0) - 3 \cdot \int_{x_0}^x P_3(s) e^{-2 \int_{x_0}^s Q_1(\tau) d\tau} ds \right] \cdot e^{-2 \int_{x_0}^x Q_1(s) ds}$$

in the view. Subordinate this solution to the condition (10) Rx

$$z(x) = \pm \left\{ \frac{1}{[y_1 - p_1(x_0)]^2} - 3 \cdot \int_{x_0}^x P_3(s) e^{-2 \int_{x_0}^s Q_1(\tau) d\tau} ds \right\}^{-\frac{1}{2}} \cdot e^{-\int_{x_0}^x Q_1(s) ds}$$

find the function. Going back to the substitution (5) above, we find the conditional solution of equation (4) in the form

$$p(x) = p_1(x) + \left\{ \frac{1}{[y_1 - p_1(x_0)]^2} - 3 \cdot \int_{x_0}^x P_3(s) e^{-2 \int_{x_0}^s Q_1(\tau) d\tau} ds \right\}^{-\frac{1}{2}} \cdot e^{-\int_{x_0}^x Q_1(s) ds},$$

and finally, we find the conditional solution of equation (1) in the form $y' = p(x)$ using the notation

$$p(x) = p_1(x) + \left\{ \frac{1}{[y_1 - p_1(x_0)]^2} - 3 \cdot \int_{x_0}^x P_3(s) e^{-2 \int_{x_0}^s Q_1(\tau) d\tau} ds \right\}^{-\frac{1}{2}} \cdot e^{-\int_{x_0}^x Q_1(s) ds}.$$

The theorem is fully proved.



REFERENCES

1. Qosimova, M. Y., & Yusupova, N. X. (2020). On a property of fractional integro-differentiation operators in the kernel of which the meyer function. *Scientific-technical journal*, 24(4), 48-50.
2. Kosimov K., Mamayusupov J. Transitions melline integral of fractional integrodifferential operators //Scientific Bulletin of Namangan State University. – 2019. – T. 1. – №. 1. – C. 12-15.
3. Azizov, M. S., & Rustamova, S. T. (2017). Yuqori tartibli differensial tenglamalarni bernulli tenglamasiga keltirib yechish. *Toshkent shahridagi turin politexnika universiteti*, 61.
4. Azizov, M., & Rustamova, S. (2019). The Task of Koshi for ordinary differential equation of first order which refer to equation of Bernoulli. *Scientific journal of the Fergana State University*, 2(1), 13-16.
5. Qosimov, H. N. (2019). Mixed fractional order integrals and derivatives which is taken multivariable function from other functionality. *Scientific Bulletin of Namangan State University*, 1(4), 3-8.
6. Tillabayev, B., & Bahodirov, N. (2021). Solving the boundary problem by the method of green's function for the simple differential equation of the second order linear. *ACADEMICIA: An International Multidisciplinary Research Journal*, 11(6), 301-304.
7. Sattorov, A. M., & Xujaxonov, Z. Z. (2019). Approach calculation of certain specific integrals by interpolating polynomials. *Scientific Bulletin of Namangan State University*, 1(3), 10-12.
8. Qosimova, M. Y., Yusupova, N. X., & Qosimova, S. T. (2021). On the uniqueness of the solution of a two-point second boundary value problem for a second-order simple differential equation solved by the bernoulli equation. *ACADEMICIA: An International Multidisciplinary Research Journal*, 11(9), 969-973.
9. Абдуразаков, А., Махмудова, Н., & Мирзамахмудова, Н. (2020). Численное решение методом прямых интеграла дифференцирования уравнений, связанных с задачами фильтрации газа. *Universum: технические науки*, (7-1 (76)), 32-35.
10. Abdurazakov, A., Makhmudova, N., & Mirzamakhmudova, N. The numerical solution by the method of direct integrals of differentiation of equations have an application in the gas filtration theorem.



11. Kosimov, L., & Kosimova, S. (2021). Optimization of the composition of dry slag-alkaline mixtures. *Збірник наукових праць Логоос*.
12. Hayotov, A. R., Bozarov, B. I., & Abduganiev, A. (2018). Optimal formula for numerical integration on two dimensional sphere. *Uzbek Mathematical Journal*, 3, 80-89.